

2022

## PHYSICS — HONOURS

Paper : CC-13

(Syllabus : 2019-2020)


[Digital Systems and Applications]

Full Marks : 50

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.*Answer *question no. 1* and *any four* from the rest.1. Answer *any five* questions :

2×5

- (a) Convert  $45.625$  into its binary equivalent.
- (b) Subtract  $(1011)_2$  from  $(1101)_2$  using 2's complement method.
- (c) Determine the output expression for the following circuit and simplify it.




- (d) Implement the boolean expression  $X = AB + \bar{A}C$  using NAND gates only.
- (e) What is the basic difference between a S-R Flip-Flop and J-K Flip-Flop?
- (f) Design a NOT-gate using a transistor ( $\beta_{\text{sat}} = 50$ ) considering  $V_{\text{CE sat}} = 0.2\text{V}$ ,  $I_C = 5\text{ mA}$  and source voltage 5 volt.
- (g) What is inequality detector?
2. (a) Simplify the following Boolean expression in SOP form using Karnaugh Map.  

$$F(A, B, C, D) = \sum m(0, 1, 2, 5, 8, 9, 10)$$
- (b) Implement the above simplified expression using basic gates.
- (c) Make the truth table for the logical function  

$$f = AB + A\bar{C} + C + AD + A\bar{B}C + ABC$$

Please Turn Over

(d) Write down the Boolean expression for the output ( $Y$ ) of the following circuit.



3+2+3+2

3. (a) Design a 8 : 1 multiplexer using two 4 : 1 multiplexers.
- (b) Implement the following Boolean expression using 8 : 1 multiplexer.

$$F(A, B, C) = \sum m(2, 4, 6, 7)$$

How can you use a 8 : 1 multiplexer to implement a logical expression with four inputs?

- (c) Write down the basic difference between decoder and de-multiplexer. 3+(3+2)+2
4. (a) Draw the circuit diagram of J-K Flip-Flop and explain its operation using sequence table.
- (b) Implement a D-Flip-Flop using J-K Flip-Flop.
- (c) Draw the full adder circuit using NAND gate only. (2+3)+2+3
5. (a) What is the basic difference in operation between MS-JK and JK Flip-Flop? Explain with block diagram.
- (b) Why is J-K Flip-Flop called an one-bit register? Explain the utility of preset and clear operation in Flip-Flop in this regard.
- (c) What is the difference between positive and negative edge triggering? Which type of triggering can be implemented using these triggering? (2+2)+(2+2)+(1+1)
6. (a) What are the differences between Synchronous and Asynchronous counters?
- (b) What is shift register? Draw a circuit diagram of a 4-bit shift register.
- (c) For 4-bit data transmission, what is the time required if we use SISO and SIPO shift register? Given the duration of each of the clock pulse is 2 ms. 2+(2+3)+3
7. (a) Draw the block diagram of D/A conversion circuit.
- (b) A five-bit D/A converter produces  $V_{out} = 0.2V$  for a digital input of 00001. Find the value of  $V_{out}$  for an input 1111.
- (c) Design a Mod-10 Asynchronous counter.
- (d) Define EPROM. 2+3+4+1

(Syllabus : 2018-2019)

[Electromagnetic Theory]

Full Marks : 50

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.*Answer **question no. 1** and **any four** questions from the rest.1. Answer **any five** questions :

2×5

- (a) Show that for electromagnetic wave propagating in free space, the electric field  $\vec{E}$ , the magnetic field  $\vec{B}$  and the unit vector in the direction of propagation  $\hat{n}$  are related by  $c\vec{B} = \hat{n} \times \vec{E}$ .
- (b) The electric field component of a plane electromagnetic wave travelling in vacuum is given by  $\vec{E}(z, t) = E_0 \cos(kz - \omega t) \hat{x}$ . Calculate the Poynting vector for this wave.
- (c) A uniform volume charge density is placed inside a conductor with resistivity  $10^{-3} \Omega \text{ m}$ .

Find the time after which the charge density becomes  $\frac{1}{e}$  of the original value.[Given :  $\epsilon_0 = 8.854 \times 10^{-12} \text{ F/m}$ ]

- (d) A certain material has a complex relative permittivity given by  $\tilde{\epsilon}_r = 40 + 12i$  at 2.45 GHz. Estimate the depth over which the amplitude of the electric field inside the material falls to half of the external value.
- (e) What should be the angle of the sun above the horizon so that sunlight reflected from a still lake is plane polarized? Given, refractive index of water = 1.33.
- (f) Describe the state of polarization of the wave represented by

$$\vec{E}(z, t) = \hat{i}E_0 \cos(kz - \omega t) - \hat{j}E_0 \sin(kz - \omega t)$$

- (g) Plane polarized light passes through a double refracting crystal of thickness  $40 \mu\text{m}$  and emerges out as circularly polarized light. If the birefringence of the crystal is  $4 \times 10^{-5}$ , find the wavelength of the incident light.

2. (a) Write down wave equations for scalar potential  $\phi(\vec{r}, t)$  and vector potential  $\vec{A}(\vec{r}, t)$  in Coulomb gauge. Consider a charge and current-free region.


- (b) Derive expressions for the electric and magnetic field in free space for the vector potential  $\vec{A} = \hat{x} a \cos(kz - \omega t) + \hat{y} b \sin(kz - \omega t)$ .

- (c) A current flowing in a long solenoid with radius R is varied such that the magnetic field inside the solenoid has magnitude  $B = \beta t^2$ , where  $\beta$  is a constant. Calculate the electric field inside and outside the solenoid and hence find the displacement current density as a function of the distance  $r$  from the axis of the solenoid.

4+3+3

Please Turn Over

3. Consider a co-axial cable of negligible resistance. If this cable is inserted between a source of constant emf and some load  $R$ , a steady current  $I$  will flow down the cable. If the emf provides a constant potential difference  $V$ , then,



- (a) What is the power supplied to the cable?
- (b) If the inner and outer radii are 'a' and 'b' respectively, then write down the expressions of  $\vec{E}$  and  $\vec{B}$  in the region ( $a < r < b$ ). [Keep in mind the cylindrical geometry]
- (c) Calculate  $\vec{S}$ , the Poynting vector.
- (d) Calculate  $\int \vec{S} \cdot d\vec{a}$  over the cross-sectional area of the cable between the inner and the outer conductors and find total electromagnetic power flow. 1+(2+2)+2+(2+1)
4. (a) An electromagnetic wave is incident on the plane interface between two different media.
- (i) Show that the wave vectors of the incident, reflected and refracted waves all lie on the same plane.
- (ii) Find the relation between the angles of incidence, reflection and refraction.
- (b) The regions of space  $z < 0$  and  $z > 0$  are filled with materials having permeabilities  $2\mu_0$  and  $5\mu_0$  respectively. The magnetic field in the region  $z > 0$  is  $\vec{B}_2 = \mu_0 (75 \hat{x} + 40 \hat{z}) T$  and there is a surface current distribution  $\vec{K} = -10 \hat{y} A/m$  at  $z = 0$ .  
Find the possible magnetic field in the region  $z < 0$ .
- (c) A plane electromagnetic wave is incident normally at the boundary of two dielectrics of refractive indices  $n_1$  and  $n_2$  ( $n_1 < n_2$ ). If the transmission co-efficient is required to be 0.80, what should be the value of  $\frac{n_2}{n_1}$ ? (2+3)+3+2

5. (a) What is displacement current? Explain why and how Ampere's circuital law for steady current was modified to include displacement current.
- (b) A plane electromagnetic wave travels in free space in the negative- $z$ -direction with a propagation constant  $20 \text{ rad/m}$ , the amplitude of the magnetic field being  $\frac{40}{377} \text{ A/m}$ . At  $t = 0, z = 0$ , the magnetic field is in the negative  $y$ -direction. Give the expression for electric field.
- (c) If vector potential  $\vec{A} = \beta x \hat{i} + 2y \hat{j} - 3z \hat{k}$  satisfies the Coulomb gauge condition, what is the value of  $\beta$ ?
- (d) Protons having the same velocity  $\vec{v} = v \hat{z}$  form an infinite beam of circular cross-section with current  $I$ . Find the direction and magnitude of the Poynting vector  $\vec{S}$  outside the beam at a distance  $r$  from its axis. (1+2)+2+2+3
6. (a) Discuss the state of polarization when the  $x$  and  $y$  components of the electric field are as follows:
- (i)  $E_x = E_0 \cos(kz + \omega t), E_y = \frac{E_0}{\sqrt{2}} \cos(kz + \omega t + \pi)$
- (ii)  $E_x = E_0 \sin\left(kz - \omega t + \frac{\pi}{3}\right), E_y = E_0 \sin\left(kz - \omega t - \frac{\pi}{6}\right)$
- (b) What will be the Brewster angle for a glass slab ( $n_g = 1.5$ ) immersed in water ( $n_w = 1.33$ )?
- (c) Four perfect polarizing plates are stacked so that the axis of each is turned  $30^\circ$  clockwise with respect to the preceding plate. How much of the intensity of an unpolarized incident beam of light is transmitted by the stack?
- (d) For calcite the values of the refractive index for o-ray and e-ray are  $n_o = 1.68134$  and  $n_e = 1.49694$  respectively, for light of wavelength  $\lambda = 404.6 \text{ nm}$ . However, corresponding to  $\lambda = 706.5 \text{ nm}$  their values are  $n_o = 1.65267$  and  $n_e = 1.48359$  respectively. A calcite quarter wave plate is construction for  $\lambda = 404.6 \text{ nm}$ . If a left circularly polarized beam of  $\lambda = 706.5 \text{ nm}$  is incident on this plate, obtain the state of polarization of the emergent beam. 3+2+2+3
7. (a) Explain the phenomenon of double refraction in a uniaxial crystal on the basis of Huygen's theory.
- (b) A  $20 \text{ cm}$  length of a certain optically active solution causes right-handed rotation of  $40^\circ$  and a  $30 \text{ cm}$  length of another solution, which does not chemically react with the first solution, causes left-handed rotation of  $24^\circ$ . What will be the optical rotation produced by  $30 \text{ cm}$  length of a mixture of the above solutions in volume ratio  $1 : 2$ ?
- (c) Plane polarized light of wavelength  $550 \text{ nm}$  is incident on a quartz crystal parallel to the optic axis. Find the least thickness for which the o-ray and the e-rays combine to form plane polarized light. Given, their refractive indices are  $\mu_o = 1.5442$  and  $\mu_e = 1.5533$  respectively.
- (d) Explain why if we hold a glass plate horizontally at the level of the eye (i.e., the angle of incidence is close to  $\pi/2$ ) the plate acts like a mirror. 2+3+3+2